

AD \_\_\_\_\_  
(Leave blank)

Award Number: W81XWH-09-1-0642

TITLE: Spectroscopic Biomarkers for Breast Tumor Grading

Á  
PRINCIPAL INVESTIGATOR: Xiaofeng Xia, PhD

CONTRACTING ORGANIZATION: The Methodist Hospital Research Institute  
Houston, TX 77030

REPORT DATE: 2010

TYPE OF REPORT: Á↔^á→

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: (Check one)

- ☒ Approved for public release; distribution unlimited
- ☐ Distribution limited to U.S. Government agencies only;  
report contains proprietary information

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. <b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</b>				
1. REPORT DATE (DD-MM-YYYY) 01-12-2010		2. REPORT TYPE Final		3. DATES COVERED (From - To) 1 SEP 2009 - 30 NOV 2010
4. TITLE AND SUBTITLE  Spectroscopic biomarkers for breast tumor grading			5a. CONTRACT NUMBER	
			5b. GRANT NUMBER W81XWH-09-1-0642	
			5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)  Xiaofeng Xia  xxia@tmhs.org			5d. PROJECT NUMBER	
			5e. TASK NUMBER	
			5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  The Methodist Hospital Research Institute, Houston, TX 77030			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)  U.S. Army Medical Research and Material Command, Fort Detrick, Maryland 21702-5012			10. SPONSOR/MONITOR'S ACRONYM(S)	
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited				
13. SUPPLEMENTARY NOTES				
14. ABSTRACT  This project aims to develop a novel magnetic resonance spectroscopy (MRS) based approach to assess the aggressiveness of breast tumors, by identifying characteristic nuclear magnetic resonance signal for breast cancer stem cells (CSCs) and using it to quantify the number of CSCs in breast tumor. Stem like cancer cells were isolated from MCF-7 breast tumor by side population (SP) sorting. 1H-NMR spectrum analysis identified two characteristic peaks of 1.28 ppm and 1.96 ppm, for the SP cells but not the unsorted cells in vitro. Application of the spectral biomarkers in vivo is still unsuccessful.				
15. SUBJECT TERMS  Cancer stem cells: Rare cancer cells that are responsible for the tumor initiation and propagation Spectroscopic biomarker: Spectra peaks that are characteristic for the cells of interest				
16. SECURITY CLASSIFICATION OF:				
a. REPORT U		17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES 7	19a. NAME OF RESPONSIBLE PERSON USAMRMC
b. ABSTRACT U	c. THIS PAGE U	UU		19b. TELEPHONE NUMBER (include area code)
Standard Form 298 (Rev. 8-98) Prescribed by ANSI Std. Z39.18				

## Table of Contents

	<u>Page</u>
Introduction.....	5
Body.....	5
Key Research Accomplishments.....	7
Conclusion.....	7
References.....	8


## Introduction

Breast tumors are routinely graded according to differentiation levels in clinics. However, its application is limited because the method is invasive and not quantitative. Clinically 30%-60% of breast tumors are graded as moderately differentiated (grade 2) which is not informative for decision making [1]. Our hypothesis is that the number of cancer stem cells (CSCs) in a breast tumor can be used as an indicator for its aggressiveness. Our hypothesis is supported by increasing evidence reported in literature. Firstly, a recent study showed that poorly differentiated tumors are enriched with cells sharing characteristic with embryonic stem cells [2]. Independent studies also indicated that tumors characterized by aggressiveness feature may encompass a higher percentage of stem-like cancer cells than the less aggressive counterparts [3, 4]. Moreover, expression levels of biomarker genes for stem-like cancer cells have been observed to be reverse correlated with prognosis clinically [5, 6]. Based on the hypothesis, we aim to develop a novel magnetic resonance spectroscopy (MRS) based approach to assess the aggressiveness of breast tumors, by identifying characteristic nuclear magnetic resonance signal for breast CSCs and using it to quantify the number of CSCs in breast tumor.

## Body

### Task 1. Identify the $^1\text{H}$ -NMR spectroscopic biomarker for stem-like cancer cells

Tumor model was created by inoculating immunodeficient SCID mice with MCF-7 human breast cancer cells. Tumors were harvested and dissociated when they grew to about 1.5 cm in diameter. In order to identify  $^1\text{H}$  NMR spectral peaks that are characteristic for breast CSCs, stem-like cancer cells were first isolated by flow cytometry as  $\text{CD44}^+\text{CD24}^{\text{-low}}\text{Lineage}^-$  cells from tumor specimen, the rest of the cells were designated as regular cancer cells and used as the control. Aqueous suspensions of cells were subjected to one-dimensional  $^1\text{H}$ -NMR spectra analysis. The result spectra were compared to identify potential characteristic peaks for stem-like cancer cells. However, diverse spectra were got for different batches of cells isolated following the same cell sorting condition. In some batches we were able to identify the characteristic 1.28ppm signal for various progenitor cells [7], while in others we failed the detect the peak (**Fig. 1**).



The above result suggest that the sorted  $CD44^{+}CD24^{-/low}Lineage^{-}$  cells are heterogenous, and may represent different types of cells for different tumors. We next sought to use different methods to isolate stem like cancer cells, aiming to purify a more homogeneous cell population. We adopted the widely used side population (SP) technique [8], in which stem like cancer cells are isolated as high dye-efflux cells. Our results ( **Fig. 2** ) showed that indeed a relatively constant percentage ( $\sim 1.5\%$ ) can be isolated following Hoechst staining.





Fig. 2 SP cells (red boxed) isolated from the tumor. Cells were dissociated and stained with  $5 \mu\text{g/ml}$  Hoechst 33342 for 90 minutes at  $37^{\circ}\text{C}$ . The Hoechst 33342 dye was excited at  $357 \text{ nm}$  and its fluorescence was analyzed under dual-wavelength of  $402\text{-}446 \text{ nm}$  (blue) and  $650\text{-}670 \text{ nm}$  (red).

We next carried out one-dimensional  $^1\text{H}$ -NMR spectra analysis again on the aqueous suspensions of cells, either on SP sorted cells or unsorted cells. Our results showed that the  $1.28 \text{ ppm}$  stem cell characteristic peak can be repeatedly identified for the SP cells. A second peak of  $1.96 \text{ ppm}$  was also repeatedly identified. Both peaks were undetectable in unsorted cells (**Fig. 3**).



**Task 2.** Test the potential of the biomarker for the quantification of stem-like cancer cells in vivo

Experiment was carried out to test the possibility of detecting the characteristic signal in vivo. Breast cancer tumor models were generated using three different methods, with the aim to create tumors with various contents of cancer stem cells. In model 1, tumors were generated by injecting regular MCF-7 breast cancer cells. In model 2, SP cells were isolated from MCF-7 cells and used for injection. In model 3, stem like cancer cells were selectively enriched by propagating MCF-7 cells in suspension as mammospheres, and the mammospheres were used for mouse injection. Animals were imaged with a 3T human MRI machine with a customized coil for small animals. However we were not able to detect either the 1.28 ppm or 1.96 ppm signal. Possible reasons were analyzed in the conclusion part.

### **Key Research Accomplishments**

We were able to isolate SP cells from MCF-7 breast tumor, which are enriched with stem like cancer cells.  $^1\text{H}$ -NMR spectrum analysis identified two characteristic peaks of 1.28 ppm and 1.96 ppm, which were repeatedly detected in SP cells but not in unsorted cells in vitro.

### **Conclusion**

To identify characteristic NMR signals for stem like cancer cells, a relatively homogenous population of cells needs to be purified. We use two widely used FACS sorting methods to isolate stem like cancer cells from human tumors generated in mouse. Surprisingly, the most widely used  $\text{CD44}^+\text{CD24}^{-/\text{low}}$  Lineage $^-$  sorting generated a population that is very likely to be highly heterogeneous. And cells prepared from different tumors appeared to have different cell contents, as indicated by the highly diverse NMR spectra. Based on cellular dye-efflux function instead of a combination of cell surface markers, the SP sorting generated a more homogeneous and repeatable population. Two NMR peaks, 1.28 ppm and 1.96 ppm, were repeatedly identified for SP cells but not unsorted cells. These signals have the potential to be used as spectroscopic biomarkers for cancer stem cells.

The effort to apply the in vitro identified spectral biomarkers in vivo, however, was unsuccessful. Till now we were unable to detect either of the characteristic signals in live animals bearing xenografted tumor. The possible reasons are (1) in vivo cells have different cellular content from in vitro cultured cells, the molecules generating the characteristic signals are not indispensable for stem cells and are not expressed in vivo; (2) the ratio of cancer stem cells in a tumor is extremely low, the characteristic signal is undistinguishable from the background under current instrument sensitivity; (3) the signals detected from SP cells in vitro are not real biomarkers for CSCs, as there is no conclusive evidence that the SP population contains real CSCs, although it is suggested to be highly enriched with stem like cancer cells; (4) although our tumor models are generated from cells with different percentages of stem like cancer cells in vitro, the tumors generated do not necessarily have different stem cell contents, as it is well known that stem cells are under constant differentiation during tumor progression.

Recent report suggests that there are naturally occurring breast cancer subset highly enriched with cancer stem cells [9]. In the future it would be a better approach to compare the tumor samples naturally containing low cancer stem cell content and tumor samples naturally containing high cancer stem cell content (known as "claudin low"). Moreover, tumors generated by xenografting the claudin low tumor specimen can serve as a better model for high CSC containing tumor, therefore will greatly facilitate the in vivo validation of the biomarkers. It is also worth noting that even the signals characteristic for the claudin low sample are not generated from real CSCs, it does not affect its clinical application, as they still have the potential to be used as an indicator for the tumor aggressiveness, which is the purpose of this project.

## References

1. Sotiriou, C., et al., *Gene expression profiling in breast cancer: understanding the molecular basis of histologic grade to improve prognosis*. J Natl Cancer Inst, 2006. 98(4): p. 262-72.
2. Ben-Porath, I., et al., *An embryonic stem cell-like gene expression signature in poorly differentiated aggressive human tumors*. Nat Genet, 2008. 40(5): p. 499-507.
3. Galli, R., et al., *Isolation and characterization of tumorigenic, stem-like neural precursors from human glioblastoma*. Cancer Res, 2004. 64(19): p. 7011-21.
4. Ponti, D., et al., *Isolation and in vitro propagation of tumorigenic breast cancer cells with stem/progenitor cell properties*. Cancer Res, 2005. 65(13): p. 5506-11.
5. Beier, D., et al., *CD133 expression and cancer stem cells predict prognosis in high-grade oligodendroglial tumors*. Brain Pathol, 2008. 18(3): p. 370-7.
6. Watanabe, O., et al., *Expression of a CD44 variant and VEGF-C and the implications for lymphatic metastasis and long-term prognosis of human breast cancer*. J Exp Clin Cancer Res, 2005. 24(1): p. 75-82.
7. Manginas, L.N., et al., *Magnetic resonance spectroscopy identifies neural progenitor cells in the live human brain*. Science, 2007. 318(5852): p. 980-5.
8. Hirschmann-Jax, C., et al., *A distinct "side population" of cells with high drug efflux capacity in human tumor cells*. Proc Natl Acad Sci U S A, 2004. 101(39): p. 14228-33.
9. Hennessy, B.T., et al., *Characterization of a naturally occurring breast cancer subset enriched in epithelial-to-mesenchymal transition and stem cell characteristics*. Cancer Res, 2009. 69(10): p. 4116-24.